



# Sugar-Sweetened Beverages\* and Young Children

**Young children drink an enormous amount of sugar.** One study found that 44% of toddlers between 19 and 24 months drink a sugar-sweetened beverage at least daily,<sup>1</sup> as do 70% of 2- to 5-year-olds.<sup>2</sup>

**Sugar-sweetened beverages are a major contributor to childhood obesity.** Every additional daily serving of soda increases a child's risk for obesity by 60 percent.<sup>3</sup> Sugar-sweetened beverage consumption in childhood also increases the risk for overweight and obesity in adulthood.<sup>4</sup>

**Sugar-sweetened beverages displace healthier items from a young child's diet.** Young children who consume a lot of sugar have a poorer overall diet quality<sup>5</sup> because these empty calories (low in nutrients) replace healthier foods that provide essential nutrients like calcium, iron, and vitamin A.<sup>6,7</sup>

**Children are drinking more sugar-sweetened beverages and less milk than they used to.**<sup>8</sup> In the 1970s, children drank nearly three times more milk than sugar-sweetened beverages. Today, they consume these beverages in equal amounts. Because milk provides essential nutrients for bone development, replacing milk with soda can reduce bone mass and increase the risk for osteoporosis later in life.<sup>10</sup>

**Sugar-sweetened beverages damage children's teeth.** Soda consumption nearly doubles the risk of dental caries in children.<sup>11</sup> The acid in soda and other sugar-sweetened beverages causes erosion of tooth enamel, often after just one sip, and the sugar in these beverages provide fuel for the bacteria that cause tooth decay. Because diet beverages also contain acid, they too increase the risk for cavities.<sup>12</sup>

**Sugar-sweetened beverages often contain caffeine.** Sixty percent of sodas sold in the United States contain caffeine.<sup>13</sup> Children who consume the amount of caffeine in one 20-oz. soda can experience withdrawal symptoms like headaches and anxiety if they stop drinking soda.<sup>14</sup>

**Breast milk is best under age one; whole milk or water from 1-2 years; low-fat, nonfat milk, and water past age 2.** Experts agree that for children under the age of one, breast milk is best. If breast-milk is not provided, then infant formula is the best alternative during the first year. From the age of one to two, children should drink whole milk and water. And once children are past the age of two, they should drink low-fat or non-fat milk and water.<sup>15</sup>

**Children should not drink more than one small serving of 100% fruit juice per day.**<sup>15</sup> Fruit juice has a high concentration of sugar and calories and lacks some of the nutrients and other benefits of whole fruit. It is best to offer children fresh whole fruit instead of juice. Infants under six months of age should not be given juice at all. Children six months to six years should not drink more than 12 ounces per day.<sup>15</sup> When children drink juice it should be 100% juice, not sugar-sweetened juice drinks.

**Diet beverages are not a healthy alternative for children.** Diet sodas are detrimental to dental health, may contain caffeine, and their long-term effects are still unknown for children.<sup>5, 16</sup>

*\* NOTE: Sugar-sweetened beverages refers to all beverages with added sugars including carbonated soft drinks, juice drinks, sports drinks, flavored and enhanced waters, sweetened teas and energy drinks.*



---

## REFERENCES

- <sup>1</sup> Fox MK, Pac S, Devaney B, Jankowski L. Feeding infants and toddlers study: what foods are infants and toddlers eating? *Journal of the American Dietetic Association*. 2004;104:S22-S30.
- <sup>2</sup> Wang YC, Bleich SN, Gortmaker SL. Increasing caloric contribution from sugar-sweetened beverages and 100% fruit juices among US children and adolescents, 1988–2004. *Pediatrics*. 2008;121:e1604-e1614.
- <sup>3</sup> Ludwig DS, et al. Relationship between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet*. 2001;357:505-508.
- <sup>4</sup> Malik VS, Popkin BM, Bray GA, Despres J-P, Hu FB. Sugar-sweetened beverages, Obesity, Type 2 Diabetes Mellitus, and Cardiovascular Disease Risk. *Circulation*. 2010;121:1356-1364.
- <sup>5</sup> Ruottinen S, et al. High sucrose intake is associated with poor quality of diet and growth between 13 months and 9 years of age: the special Turku coronary risk factor intervention project. *Pediatrics*. 2008; 121:e1676-e1685.
- <sup>6</sup> Healthy Eating Research. The negative impact of sugar-sweetened beverages on children's health. Robert Wood Johnson Foundation. November 2009.
- <sup>7</sup> Kranz S, Smicklas-Wright H, Siega-Riz A, et al. Adverse Effect of High Added Sugar Consumption on Dietary Intake in American Preschoolers. *Journal of Pediatrics*, 2005; 146(1): 105–111.
- <sup>8</sup> Harnack L, Stang J, Story M. Soft drink consumption among US children and adolescents: nutritional consequences. *J Am Diet Assoc*. 1999;99:436–441.
- <sup>9</sup> Smith TA, Lin, B-H Lin, Lee J-Y. Taxing caloric sweetened beverages: potential effects on beverage consumption, calorie intake, and obesity, ERR-100, U.S. Department of Agriculture, Economic Research Service, July 2010.
- <sup>10</sup> Ondrak KS, Morgan DW. Physical activity, calcium intake and bone health in children and adolescents. *Sports Med* 2007; 37:587-600.
- <sup>11</sup> Sohn W, Burt BA, Sowers MR. Carbonated Soft Drinks and Dental Caries in the Primary Dentition. *J Dent Res*. 2006; 85(3): 262-266.
- <sup>12</sup> Buyer DM. Are you drinking your teeth away? How soda and sports drinks dissolve enamel. *Journal of the Indiana Dental Association*. 2009. Summer: 11-13.
- <sup>13</sup> Keast RSJ, Riddell LJ. Caffeine as a flavor additive in soft drinks. *Appetite*. 2007; 49:255-259.
- <sup>14</sup> Heatherley SV, Hancock KM, Rogers PJ. Psychostimulant and other effects of caffeine in 9- to 11-year-old children. *J Child Psychol Psychiatry*. 2006;47(2):135-42.
- <sup>15</sup> American Academy of Pediatrics and American Heart Association. Dietary Recommendations for Children and Adolescents: A Guide for Practitioners. *Pediatrics* 2006;117:544-559
- <sup>16</sup> Ludwig DS. Artificially Sweetened Beverages Cause for Concern. *JAMA*. 2009;302(22):2477-2478.

This fact sheet was reviewed by a scientific advisory panel of nationally recognized scientists, researchers and public health experts convened by the **California Center for Public Health Advocacy**. For more info about advisory panel, contact [info@kickthecan.info](mailto:info@kickthecan.info). | Last Updated: January 31, 2011.